



Behaviour of Oriza Stiva Under Variable UV Radiations

Asad Waseem^{1*}, Rizwan Akhtar², Amna Ali³, Adil Latif⁴

^{1*2,3,4}Department of Space Science, University of the Punjab, Lahore

* Asad Waseem Email: asadwaseem1514@hotmail.com

Citation | A. Waseem, Rizwan Akhtar, Amna Ali, and Adil Latif, “Behaviour of Oriza Stiva Under Variable UV Radiations”, IJASD, vol. 3, no. 3, pp. 57–63, Aug. 2021.

Received | July 12, 2021; **Revised** | Aug 18, 2021; **Accepted** | Aug 21, 2021;

Published | Aug 26, 2021

Solar UV radiation at the earth's surface grows as the ozone layer depletes, affecting economically vital plants. The effects of UV radiation on germination, growth, chlorophyll content, and fresh and dry weight of Oriza Stiva were investigated in this study. Oriza Stiva seeds were placed in plates and subjected to UV light for 30, 60, and 120 minutes each day. UV radiation lowered the percentage of seeds that germinated. Increased UV exposure significantly slowed seedling root and shoot growth. In comparison to the control, UV irradiation had a beneficial effect on chlorophyll contents, increasing total chlorophyll content. According to the findings, UV radiation reduced radicle and plumule lengths while increasing chlorophyll content in Oriza Stiva.

Keywords: Germination, Ozone, Oriza Stiva.

Introduction

Plants rely on solar energy for their growth and development. A small variation in light wavelength can greatly impact plant physiology and biochemistry [1][2]. Light waves with wavelengths of 400-700 nm have a stronger impact on plant responses, causing changes in the magnitude relationship of UV light to influence biomass allocation [3][4][5]. The presence of the ozone layer at around 10-30 km above the earth's surface protects humans from damaging UV rays all across the world[6][2]. The depleted stratospheric layer and global weather change are two of the most serious risks. UV light with a 280-320 nm wavelength can reach the earth's surface if the ozone layer is depleted. As the ozone layer around the globe thins, the harmful effects of UV light rise, and their strength is amplified by a variety of topographic conditions[7]. In 1881, Hartley first observed the negative effects of UV light [8][9], [10].

UV light has a greater impact at high altitudes than in low-lying places, and it affects practically all plant activities beginning with germination. The ozone layer's depletion increased the penetration of solar UV radiation to the earth's surface. UV radiation that reaches the earth's surface can impact plant growth. The plumule and radicle length of freshly emerging plants are similarly affected by daily UV shocks [11]. UV light of high intensity is harmful to all creatures, including microorganisms [12]–[14]. Paul (2001) investigated species and discovered that UV exposure can produce physiological, morphological, and chemical changes as well as molecular changes. UV radiation reduces the formation of chlorophyll, soluble proteins, and Rubisco in C3 plants [15][16]. UV light inhibits development, destroys photosynthetic pigments, lowers carbon intake, and affects various other functional and hormonal systems as a stressor [17], [18][19]. Plants experience intraspecific differences in biochemicals, wet or dry mass, and other factors due to UV radiation [20][21]. Most plant species are sensitive to UV light, whereas only a few are unaffected [22][23]. UV radiation has a negative impact on physiological and metabolic systems. UV-treated seeds are disrupted by protein formation, gas exchange, water transport, and hormonal and enzymatic activity[24]–[26]. UV therapy is directly or indirectly detrimental to plants [27], [28]. It has been found that UV therapy reduces photosystem II activity in photosynthesis. UV radiation causes the stomatal opening and shutting to be disrupted, inhibiting gaseous exchange and lowering transpiration rates [29]–[31].

UV light causes changes in cell division, hormone abnormalities, and nucleic acid damage [32][33]. UV light produces curled leaves and a reduction in plant size, leaf size, and biomass[34][35]. *Oriza Stiva* members are utilized as potherbs and are essential for their chemical components, flavonoids, and mineral elements. UV radiation reduced the quantity of chlorophyll, anthocyanin, and proline in vegetable plants, according to Mahdavian[36][37].

The goal of this study was to see how UV radiation influences seed germination. Specifically, in plants that are acclimating to cool habitats and are thought to be vulnerable to high temperatures and in plants that have been exposed to UV light. Seeds of various *Oriza Stiva* species were utilized to screen for UV sensitivity in this investigation.

Materials and Procedures

Because of the importance of these plants as pot herbs, seeds of *Oryza Sativa* were utilized in the experiment. These species' seeds were harvested in good health. Clean seeds of *Oryza Sativa* were sterilized for two minutes with 0.5 percent sodium hypochlorite, then rinsed and steeped for two hours in distilled water. The soaking 20 seeds were placed in Petri plates with two discs of Whatman no.1 filter paper. Seeds were divided into 9 groups for treatment, each with three replicates. One group was for control, two groups for 25 minutes, two groups for 55 minutes, and two groups for 120 minutes. The six groups were exposed to UV radiation for 25, 55, and 120 minutes every day.

There were six treatments, each with three replicates. Petri plates were kept at room temperature for the germination investigation. Daily observations on germination were kept

track of. To maintain the Petri plates moist, little volumes of distilled water were introduced on a regular basis. Germination was tracked on a daily basis for up to 20 days. The pigments of chlorophyll were determined on the 14th day. The seedlings' radicle and plumule lengths and their fresh and dry weights were measured at the end of the experiment. On the 16th day of the experiment, the plumule and radical lengths were measured [38].

The fresh weight of each plate was determined on the 16th day of seed germination, and the plants were subsequently dried in the oven for 48 hours. The dry weight was determined the next day.

Two-hour soaking seeds were exposed to UV light for 0 (control), and chlorophyll in the leaf was measured using a chlorophyll meter on the 12th day.

Results

UV light at high intensities causes a variety of changes, both outwardly and inside, and chemical disruption [39]. With more UV radiation exposure, the overall germination rate was dramatically lowered compared to the control group. The irradiation rate of ultraviolet radiation is reduced and it increases the germination percentage. This investigation revealed that UV irradiation had a significant impact on seed germination and chlorophyll content in the plants studied. UV radiation treated rice for 25, 55, and 120 minutes, respectively.

These findings were consistent with the existing literature. Seeds exposed to UV light during germination and seedling appearance have a greater impact than seeds exposed to UV light at later stages [40].

Lengths of radicle and plumule

Oryza sativa showed a similar reduction in shoot length when compared to the control group. Seeds and seedlings in their early phases of development are extremely sensitive to environmental factors, particularly UV light [41], [42]. The plant growth stage and environmental conditions influence UV-irradiated seeds germination and chlorophyll concentration [43], [44]. Previous research has revealed that not only germination percentage, but also shoot length, is influenced in particular plants [45]. Phenols are induced to accumulate in plants during exposure to UV radiation as a stress signal [46], [47] and protect plants from stressors [48]–[51].

The effects of UV radiation on germination and growth are determined by the fresh and dry weight. The fresh weight was determined on the 16th day after germination. Compared to the Control, both the Fresh and Dry weights were reduced. The fresh and dry weight in control replicates is higher than in UV-treated replicates. Flavonoids, found in seed coats, protect seeds [52]. The current study found that shoot and root length were affected in all treatments after being exposed to UV radiation. UV radiation had an effect on the root and shoot lengths of sprouting seedlings in this investigation. Exposure to UV light suppressed root and shoot growth, according to the author [52]. Similarly, the fresh and dry weight of the seedling, which indicates seedling growth, decreases. According to Mladin UV light damages photosystem II, resulting in reduced or even no photosynthesis and low biomass. Plants are affected by UV light, although their defense mechanisms try to mitigate this by activating their antioxidant system [53]. UV light increases antioxidant enzyme activity. UV light also increases the activity of antioxidant enzymes [54].

The fresh and dry weight determines UV light's impact on germination and growth. The fresh. Control duplicates in both species have higher fresh and dry weights than UV-treated replicates. Seed coverings include flavonoids, which preserve seeds. In the current study, the shoot and root length of emerging seedlings. According to research UV radiation inhibited root and shoot growth, and the seedling's fresh and dry weight, which reflects seedling growth, also drops. UV light affects photosystem II, leading in reduce or no photosynthesis and low biomass, according to Mladin et al. (2014). UV light affects plants, although their defense mechanisms try to counteract this by activating their antioxidant

system. UV radiation boosts the activity of antioxidant enzymes. According to Kondo and Kawashima, UV radiation also boosts the activity of antioxidant enzymes (2000).

Conclusions

Seeds exposed to UV radiation had a significantly lower germination percentage and a slower germination rate. The radicle and plumule lengths were altered by UV exposure. In all UV exposures, shoot length was reduced compared to the control, but root length remained unchanged. UV exposure also caused seedlings to curl and twist. UV radiation lowered the fresh and dry weight of the plant; in comparison to the control, UV exposure reduced the fresh and dry weight substantially. UV radiation can also enhance the number of chlorophyll pigments in the plant compared to controls. Plant development, seedling, and germination were slowed by UV radiation, although chlorophyll content rose.

References

- [1] Y. Roupahel, M. C. Kyriacou, S. A. Petropoulos, S. De Pascale, and G. Colla, "Scientia Horticulturae Improving vegetable quality in controlled environments," *Sci. Hortic. (Amsterdam)*, vol. 234, no. February, pp. 275–289, 2018, doi: 10.1016/j.scienta.2018.02.033.
- [2] J. Schmitt and R. D. Wulff, "Light spectral quality, phytochrome and plant competition," *Trends Ecol. Evol.*, vol. 8, no. 2, pp. 47–51, Feb. 1993, doi: 10.1016/0169-5347(93)90157-K.
- [3] P. Taylor, "Critical Reviews in Plant Sciences UV-B-Induced Secondary Plant Metabolites - Potential Benefits for Plant and Human Health UV-B-Induced Secondary Plant Metabolites - Potential Benefits for Plant and Human Health," no. February 2013, pp. 37–41, 2012, doi: 10.1080/07352689.2012.664979.
- [4] A. A. HaniaArif, Mamoona Midhat Kazmi, Anam Munwar, "Improving Rice Yield Through Insufficient Water," *Int. J. Agric. Sustain. Dev.*, vol. 3, no. 2, pp. 40–45, 2021.
- [5] S. K. Maliakal, K. McDonnell, S. A. Dudley, and J. Schmitt, "Effects of Red to Far-Red Ratio and Plant Density on Biomass Allocation and Gas Exchange in *Impatiens capensis*," <https://doi.org/10.1086/314157>, vol. 160, no. 4, pp. 723–733, Jul. 2015, doi: 10.1086/314157.
- [6] A. Khaleghi, R. Naderi, C. Brunetti, and B. E. Maserti, "Morphological, physiochemical and antioxidant responses of *Maclura pomifera* to drought stress," *Sci. Rep.*, pp. 1–12, 2019, doi: 10.1038/s41598-019-55889-y.
- [7] K. R. Reddy, V. G. Kakani, D. Zhao, S. Koti, and W. Gao, "Interactive Effects of Ultraviolet-B Radiation and Temperature on Cotton Physiology, Growth, Development and Hyperspectral Reflectance," <https://doi.org/10.1562/2003-11-19-RA.1>, vol. 79, no. 5, pp. 416–427, May 2004, doi: 10.1562/2003-11-19-RA.1.
- [8] C. Jiao, R. Yang, Y. Zhou, and Z. Gu, "Nitric oxide mediates isoflavone accumulation and the antioxidant system enhancement in soybean sprouts," *FOOD Chem.*, 2016, doi: 10.1016/j.foodchem.2016.02.147.
- [9] W. N. Hartley, "On the absorption of solar rays by atmospheric ozone," *J. Chem. Soc. Trans.*, vol. 39, no. 0, pp. 111–128, Jan. 1881, doi: 10.1039/CT8813900111.
- [10] M. Tevini and A. H. Teramura, "UV-B EFFECTS ON TERRESTRIAL PLANTS," *Photochem. Photobiol.*, vol. 50, no. 4, pp. 479–487, Oct. 1989, doi: 10.1111/J.1751-1097.1989.TB05552.X.
- [11] S. D. Flint and M. M. Caldwell, "A biological spectral weighting function for ozone depletion research with higher plants," *Physiol. Plant.*, vol. 117, no. 1, pp. 137–144, Jan. 2003, doi: 10.1034/J.1399-3054.2003.1170117.X.
- [12] S. U. Zaman, K. Ali, W. Khan, M. Ali, T. Jan, and M. Nisar, "Ethno-botanical and geo-referenced profiling of medicinal plants of Nawagai Valley, District Buner (Pakistan)," *Biosyst. Divers.*, vol. 26, no. 1, pp. 56–61, 2018, doi: 10.15421/011809.

- [13] P. W. Barnes, S. Maggard, S. R. Holman, and B. S. Vergara, "Intraspecific Variation in Sensitivity to UV-B Radiation in Rice," *Crop Sci.*, vol. 33, no. 5, pp. 1041–1046, Sep. 1993, doi: 10.2135/CROPSCI1993.0011183X003300050033X.
- [14] E. Goto, K. Hayashi, S. Furuyama, S. Hikosaka, and Y. Ishigami, "Effect of UV light on phytochemical accumulation and expression of anthocyanin biosynthesis genes in red leaf lettuce," pp. 179–186, 2016, doi: 10.17660/ActaHortic.2016.1134.24.
- [15] C. Inostroza-blancheteau, P. Acevedo, R. Loyola, P. Arce-johnson, M. Alberdi, and M. Reyes-díaz, "Plant Physiology and Biochemistry Short-term UV-B radiation affects photosynthetic performance and antioxidant gene expression in highbush blueberry leaves," *Plant Physiol. Biochem.*, vol. 107, pp. 301–309, 2016, doi: 10.1016/j.plaphy.2016.06.019.
- [16] L. E.-A. A. Nasser, "Effects of UV-B radiation on some physiological and biochemical aspects in two cultivars of barley (*Hordeum vulgare* L.)," *Egypt. J. Biol.*, vol. 3, no. 1, Jul. 2004, doi: 10.4314/ejb.v3i1.29924.
- [17] A. Fukushima and K. Saito, "Influence of UV-light on carthamin accumulation and floret elongation in a cultivar of saffron thistle (*Carthamus tinctorius* L.)," *Acta Physiol. Plant.* 2000 222, vol. 22, no. 2, pp. 159–162, 2000, doi: 10.1007/S11738-000-0071-9.
- [18] F. Hollósy, "Effects of ultraviolet radiation on plant cells," *Micron*, vol. 33, no. 2, pp. 179–197, Jan. 2002, doi: 10.1016/S0968-4328(01)00011-7.
- [19] T. M. Asad waseem, Aamer Amin, Jamal Hassan, "Impact Assessment of Forest Managment on Inhabitants," *Int. J. Agric. Sustain. Dev.*, vol. 3, no. 2, pp. 31–39, 2021.
- [20] H. Dou and G. Niu, "Pre-Harvest UV-B Radiation and Photosynthetic Photon Flux Density Interactively Affect Plant Photosynthesis, Growth, and Secondary Metabolites Accumulation in Basil (*Ocimum basilicum*) Plants," 2019.
- [21] C. A. Kolb, M. A. Käser, J. Kopecký, G. Zotz, M. Riederer, and E. E. Pfündel, "Effects of Natural Intensities of Visible and Ultraviolet Radiation on Epidermal Ultraviolet Screening and Photosynthesis in Grape Leaves," *Plant Physiol.*, vol. 127, no. 3, pp. 863–875, Nov. 2001, doi: 10.1104/PP.010373.
- [22] Z. Guadalupe and M. P. Diago, "Phenolic composition of Tempranillo wines following early defoliation of the vines and Javier Tardaguila," no. August 2011, pp. 925–934, 2012, doi: 10.1002/jsfa.4671.
- [23] A. H. Teramura, "Effects of ultraviolet-B radiation on the growth and yield of crop plants," *Physiol. Plant.*, vol. 58, no. 3, pp. 415–427, Jul. 1983, doi: 10.1111/J.1399-3054.1983.TB04203.X.
- [24] W. Khan, M. Hussain, K. Ali, M. Ali, and M. Nisar, "Distribution and phenotypic variation in *Juglans regia* L. growing in Hindu Kush ranges of Pakistan," *Acta Ecol. Sin.*, vol. 40, no. 5, pp. 363–372, Oct. 2020, doi: 10.1016/J.CHNAES.2020.02.009.
- [25] B. Hieng, K. Ugrinović, J. Šuštar-Vozlič, and M. Kidrič, "Different classes of proteases are involved in the response to drought of *Phaseolus vulgaris* L. cultivars differing in sensitivity," *J. Plant Physiol.*, vol. 161, no. 5, pp. 519–530, Jan. 2004, doi: 10.1078/0176-1617-00956.
- [26] C. Pastore, S. Zenoni, M. Fasoli, M. Pezzotti, G. B. Tornielli, and I. Filippetti, "Selective defoliation affects plant growth, fruit transcriptional ripening program and flavonoid metabolism in grapevine," 2013.
- [27] G. Horneck, U. Eschweiler, G. Reitz, J. Wehner, R. Willimek, and K. Strauch, "Biological responses to space: Results of the experiment 'Exobiological Unit' of ERA on EURECA I," *Adv. Sp. Res.*, vol. 16, no. 8, pp. 105–118, Jan. 1995, doi: 10.1016/0273-1177(95)00279-N.
- [28] P. N. Joshi et al., "Partial protection of photosynthetic apparatus from UV-B-induced

- damage by UV-A radiation,” *Environ. Exp. Bot.*, vol. 59, no. 2, pp. 166–172, Mar. 2007, doi: 10.1016/J.ENVEXPBOT.2005.11.005.
- [29] T. A. Day and T. C. Vogelmann, “Alterations in photosynthesis and pigment distributions in pea leaves following UV-B exposure,” *Physiol. Plant.*, vol. 94, no. 3, pp. 433–440, Jul. 1995, doi: 10.1111/J.1399-3054.1995.TB00950.X.
- [30] U. C. Shukla, P. C. Joshi, and P. Kakkar, “Synergistic Action of Ultraviolet-B Radiation and Cadmium on the Growth of Wheat Seedlings,” *Ecotoxicol. Environ. Saf.*, vol. 51, no. 2, pp. 90–96, Feb. 2002, doi: 10.1006/EESA.2001.2136.
- [31] J. Jiao, Q. Gai, W. Wang, M. Luo, C. Gu, and Y. Fu, “Ultraviolet Radiation-Elicited Enhancement of Iso flavonoid,” 2015, doi: 10.1021/acs.jafc.5b03138.
- [32] I. U. U. Radiation, “Impact of Increasing Ultraviolet-B (UV-B) Radiation on Photosynthetic Processes,” *J. Photochem. Photobiol. B Biol.*, 2014, doi: 10.1016/j.jphotobiol.2014.02.004.
- [33] S. W. Mpoloka, “Effects of prolonged UV-B exposure in plants,” *African J. Biotechnol.*, vol. 7, no. 25, pp. 4874–4883, 2008.
- [34] N. H. Furness, M. K. Upadhyaya, and D. P. Ormrod, “Seedling growth and leaf surface morphological responses of three rangeland weeds to ultraviolet-B radiation,” *Weed Sci.*, vol. 47, no. 4, pp. 427–434, 1999, doi: 10.1017/S0043174500092031.
- [35] A. A. Saira Batool , Maryam M. Ali , Zainab Tahir, “Investigation of Characteristics of Hydrological Droughts in Indus Basin,” *Int. J. Agric. Sustain. Dev.*, vol. 3, no. 2, pp. 46–53, 2021.
- [36] K. Mahdavian, M. Ghorbanli, and K. M. Kalantari, “The effects of ultraviolet radiation on some antioxidant compounds and enzymes in *Capsicum annum* L,” *Turk. J. Botany*, vol. 32, no. 2, pp. 129–134, 2008.
- [37] H. Frohnmeyer and D. Staiger, “Update on Ultraviolet-B Light Responses Ultraviolet-B Radiation-Mediated Responses in Plants . Balancing Damage and Protection 1,” vol. 133, no. December, pp. 1420–1428, 2003, doi: 10.1104/pp.103.030049.deeper.
- [38] A. A. Bushra Noreen, Namra Ghaffar, Maryam Muhammad Ali, “Rice Yeild Estimation in Sawat Region Incorporating The Local Physio-Climatic Parameters,” *Int. J. Agric. Sustain. Dev.*, vol. 3, no. 2, pp. 54–59, 2021.
- [39] S. S. Shaukat, M. A. Farooq, M. F. Siddiqui, and S. Zaidi, “Effect of enhanced UV-B radiation on germination, seedling growth and biochemical responses of *Vigna mungo* (L.) Hepper,” *Pakistan J. Bot.*, vol. 45, no. 3, pp. 779–785, 2013.
- [40] M. Saile-Mark and M. Tevini, “Effects of solar UV-B radiation on growth, flowering and yield of central and southern European bush bean cultivars (L.,” *UV-B Biosph.*, pp. 114–125, 1997, doi: 10.1007/978-94-011-5718-6_11.
- [41] S. E. Sultan, “Phenotypic plasticity for plant development, function and life history,” *Trends Plant Sci.*, vol. 5, no. 12, pp. 537–542, Dec. 2000, doi: 10.1016/S1360-1385(00)01797-0.
- [42] A. Pliura, A. Baliuckiene, and V. Baliuckas, “Phenogenetic response of silver birch populations and half-sib families to elevated ozone and ultraviolet-B radiation at juvenile age,” *Environ. Pollut.*, vol. 156, no. 1, pp. 152–161, Nov. 2008, doi: 10.1016/J.ENVPOL.2007.12.023.
- [43] K. Ali et al., “The ethnobotanical domain of the Swat Valley, Pakistan,” *J. Ethnobiol. Ethnomed.*, vol. 14, no. 1, pp. 1–15, Jun. 2018, doi: 10.1186/S13002-018-0237-4/TABLES/6.
- [44] F. Kuhlmann and C. Müller, “Development-dependent effects of UV radiation exposure on broccoli plants and interactions with herbivorous insects,” *Environ. Exp. Bot.*, vol. 66, no. 1, pp. 61–68, Apr. 2009, doi: 10.1016/J.ENVEXPBOT.2009.01.001.
- [45] B. Peykarestan, M. Seify, M. S. Fadaei, and M. Hatim, “UV irradiation effects on seed

- germination and growth, protein content, peroxidase and protease activity in *Portulaca grandiflora* and *Portulaca oleracea*,” *World Appl. Sci. J.*, vol. 17, no. 7, pp. 802–808, 2012.
- [46] M. Kozłowska, E. Brzezińska, and M. Stobiecki, “Sensitivity differences and accumulation of screening compounds in three conifer plants under enhanced UV-B radiation,” *Polish J. Environ. Stud.*, vol. 16, no. 6, pp. 823–830, 2007.
- [47] K. Ravindran, A. Indrajith, V. Balakrishnan, K. Venkatesan, and G. Kulandaivelu, “Determination of defense mechanism in *Phaseolustrilobus* Ait.: Seedlings treated under UV-B radiation,” *African Crop Sci. J.*, vol. 16, no. 2, pp. 111–118, 2010, doi: 10.4314/acsj.v16i2.54351.
- [48] D. C. Close and C. McArthur, “Rethinking the role of many plant phenolics – protection from photodamage not herbivores?,” *Oikos*, vol. 99, no. 1, pp. 166–172, Oct. 2002, doi: 10.1034/J.1600-0706.2002.990117.X.
- [49] I. N. De Abreu and P. Mazzafera, “Effect of water and temperature stress on the content of active constituents of *Hypericum brasiliense* Choisy,” *Plant Physiol. Biochem.*, vol. 43, no. 3, pp. 241–248, Mar. 2005, doi: 10.1016/J.PLAPHY.2005.01.020.
- [50] N. A. Olenichenko and N. V. Zagoskina, “Response of Winter Wheat to Cold: Production of Phenolic Compounds and L-Phenylalanine Ammonia Lyase Activity,” *Appl. Biochem. Microbiol.* 2005 416, vol. 41, no. 6, pp. 600–603, Nov. 2005, doi: 10.1007/S10438-005-0109-2.
- [51] G. Ganeva and E. Zozikova, “Effect of increasing Cu Concentration on Growth and Content of Free Phenols in Two Lines of Wheat (*Triticum Aestivum*) With Difference Tolerance,” *Gen. Appl. Plant Physiol.*, vol. 33, no. May, pp. 75–82, 2007.
- [52] S. S. Shaukat, S. Zaidi, and M. A. Khan, “Effect of Supplemental Uv-B Radiation on Germination , Seedling Growth , and Biochemical Responses of Sunflower (*Helianthus Annuus* L .),” *FUUAST J. Biol.*, vol. 1, no. 1, pp. 27–33, 2011.
- [53] E. Horváth, G. Szalai, and T. Janda, “Induction of Abiotic Stress Tolerance by Salicylic Acid Signaling,” *J. Plant Growth Regul.* 2007 263, vol. 26, no. 3, pp. 290–300, Aug. 2007, doi: 10.1007/S00344-007-9017-4.
- [54] N. Kondo and M. Kawashima, “Enhancement of the Tolerance to Oxidative Stress in Cucumber (*Cucumis sativus* L.) Seedlings by UV-B Irradiation: Possible Involvement of Phenolic Compounds and Antioxidative Enzymes,” *J. Plant Res.* 2000 1133, vol. 113, no. 3, pp. 311–317, 2000, doi: 10.1007/PL00013863.



Copyright © by authors and 50Sea. This work is licensed under Creative Commons Attribution 4.0 International License.